




We claim:


1. A method of treating a neurological activity in an animal, comprising:
administering to said animal an effective amount of a compound having an
affinity for FKBP-type immunophilins according to formula I



Formula I

or a pharmaceutically acceptable salt thereof,

- wherein Y is CH₂, O, NH, or N-(C1-C4 alkyl);
- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns, or



- wherein Q is hydrogen, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl;
- wherein T is Ar or substituted 5-7 membered cycloalkyl with substituents at positions 3 and 4 which are independently selected from the group consisting of hydrogen, hydroxyl, O-(C1-C4)-alkyl or O-(C1-C4)-alkenyl and carbonyl;
- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, 1,2-methylenedioxy, carbonyl and phenyl;
- wherein R₁ is either hydrogen or U; X is either oxygen or CH-U, provided that if R₁ is hydrogen, then X is CH-U, or if X is oxygen then R₁ is U;
- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar (Ar as described above);
- wherein J is hydrogen or C1 or C2 alkyl or benzyl; K is (C1-C4)-straight or branched alkyl, benzyl or cyclohexylethyl; or wherein J and K may be taken

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

together to form a 5-7 membered heterocyclic ring which may contain an oxygen (O), sulfur (S), SO or SO₂ substituted therein;

D1

- wherein n is 0-3; and

- wherein said neurological activity does not include amyotrophic lateral sclerosis.

2. The method of claim 1, wherein the neurological activity is selected from the group consisting of stimulation of damaged neurons, promotion of neuronal regeneration, prevention of neurodegeneration, and treatment of a neurological disorder.

3. The method of claim 2, wherein the neurological disorder is selected from the group consisting of peripheral neuropathies cause by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, stroke associated with brain damage, and neurological disorders relating to neurodegeneration.

4. The method of claim 3, wherein the neurological disorder is Alzheimer's Disease or Parkinson's Disease.

5. The method of claim 1, wherein J and K are taken together to form a 5 membered heterocyclic ring.

6. The method of claim 5,

- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-

D2

straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns;

- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar; and

- wherein Ar is selected from the group consisting monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, and 1,2-methylenedioxy.

7. The method of claim 6, wherein

- if X is O, Y is O, NH or N-(C1-C4 alkyl), R₁ is C1-C6 straight or branched alkyl, C2-C6 straight or branched alkenyl, C5-C7 cycloalkyl or cycloalkenyl substituted with C1-C4 straight or branched alkyl or C2-C4 straight or branched alkenyl, and Ar is 1-naphthyl, 2-naphthyl, indolyl, 2-furyl, 3-furyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl

- then (CH₂)_nZ and R₂ taken together do not form:

- 1) substituted or unsubstituted indolyl, 2-furyl, 3-furyl, thiazolyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, phenyl
- 2) an alkyl or alkenyl chain with substituted or unsubstituted indolyl, 2-furyl, 3-furyl, thiazolyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, phenyl
- 3) an alkyl or alkenyl chain substituted with C5-C7 cycloalkyl.

8. The method of claim 1, wherein J and K are taken together to form a 6 membered heterocyclic ring.

9. The method of claim 8,

- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns;

- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar; and
- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, and 1,2-methylenedioxy.

10. The method of claim 9, wherein
 - if n is 0, X is O, Y is O, R₁ is 1,1-dimethyl-1-propyl, and R₂ is 2-phenylethyl, then Z is not 1,1-dimethyl-1-prop-2-enyl, phenyl, cyclohexyl, or 3-(N,N-diallyl)benzamide; and
 - if n is 0, X is O, Y is O, R₁ is tert-butyl, and R₂ is 2-(3',4',5'-trimethoxyphenyl)ethyl, then Z is not phenyl.
11. The method according to claim 1, further comprising co-administering to said

animal an effective amount of a neurotrophic factor selected from the group consisting of nerve growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3.

12. The method of claim 11, wherein the neurological activity is selected from the group consisting of stimulation of damaged neurons, promotion of neuronal regeneration, prevention of neurodegeneration and treatment of a neurological disorder.

13. The method of claim 12, wherein the neurological disorder is selected from the group consisting of peripheral neuropathies caused by physical injury or disease state, physical damage to the brain, physical damage to the spinal cord, and neurological disorders relating to neurodegeneration.


14. The method of claim 12, wherein the neurological disorder is Alzheimer's Disease or Parkinson's Disease.

15. The method of claim 11, wherein J and K are taken together to form a 5 membered heterocyclic ring.

16. The method of claim 11, wherein J and K are taken together to form a 6 membered heterocyclic ring.

17. A method for preventing neurodegeneration in an animal, comprising:


administering to said animal an effective amount of a compound having an affinity for FKBP-type immunophilins according to formula I



Formula I

or a pharmaceutically acceptable salt thereof,

- wherein Y is CH₂, O, NH, or N-(C1-C4 alkyl);
- wherein X and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns, or



- wherein Q is hydrogen, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl;

- wherein T is Ar or substituted 5-7 membered cycloalkyl with substituents at positions 3 and 4 which are independently selected from the group consisting of hydrogen, hydroxyl, O-(C1-C4)-alkyl or O-(C1-C4)-alkenyl and carbonyl;
- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, 1,2-methylenedioxy, carbonyl and phenyl;
- wherein R₁ is either hydrogen or U; X is either oxygen or CH-U, provided that if R₁ is hydrogen, then X is CH-U, or if X is oxygen then R₁ is U;
- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar (Ar as described above);
- wherein J is hydrogen or C1 or C2 alkyl or benzyl; K is (C1-C4)-straight or branched alkyl, benzyl or cyclohexylethyl; or wherein J and K may be taken together to form a 5-7 membered heterocyclic ring which may contain an oxygen (O), sulfur (S), SO or SO₂ substituted therein;

- wherein n is 0-3; and
- wherein the neuronal degeneration is not caused by amyotrophic lateral sclerosis.

18. The method of claim 17, wherein J and K are taken together to form a 5 membered heterocyclic ring.

19. The method of claim 18,

- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns;
- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar; and
- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently

selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, and 1,2-methylenedioxy.

20. The method of claim 19, wherein

- if X is O, Y is O, NH or N-(C1-C4 alkyl), R₁ is C1-C6 straight or branched alkyl, C2-C6 straight or branched alkenyl, C5-C7 cycloalkyl or cycloalkenyl substituted with C1-C4 straight or branched alkyl or C2-C4 straight or branched alkenyl, and Ar is 1-naphthyl, 2-naphthyl, indolyl, 2-furyl, 3-furyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl

- then (CH₂)_nZ and R₂ taken together do not form:

- 1) substituted or unsubstituted indolyl, 2-furyl, 3-furyl, thiazolyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, phenyl
- 2) an alkyl or alkenyl chain with substituted or unsubstituted indolyl, 2-furyl, 3-furyl, thiazolyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, phenyl
- 3) an alkyl or alkenyl chain substituted with C5-C7 cycloalkyl.

21. The method of claim 17, wherein J and K are taken together to form a 6 membered heterocyclic ring.

- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns;

- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar; and

- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, and 1,2-methylenedioxy.

23. The method of claim 22, wherein


-if n is 0, X is O, Y is O, R₁ is 1,1-dimethyl-1-propyl, and R₂ is 2-phenylethyl, then Z is not 1,1-dimethyl-1-prop-2-enyl, phenyl, cyclohexyl, or 3-(N,N-diallyl)benzamide; and

-if n is 0, X is O, Y is O, R₁ is tert-butyl, and R₂ is 2-(3',4',5'-trimethoxyphenyl)ethyl, then ^DZ is not phenyl.

24. The method of claim 17, further comprising co-administering an effective amount of a neurotrophic factor to prevent neurodegeneration selected from the group consisting of nerve growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, and neurotrophin-3.

25. A method for promoting neuronal regeneration and growth in animals, comprising:


administering to said animal an effective amount of compound having an affinity for FKBP-type immunophilins according to formula I



Formula I

or a pharmaceutically acceptable salt thereof,

- wherein Y is CH₂, O, NH, or N-(C1-C4 alkyl);
- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns, or



23

- wherein Q is hydrogen, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl;
- wherein T is Ar or substituted 5-7 membered cycloalkyl with substituents at positions 3 and 4 which are independently selected from the group consisting of hydrogen, hydroxyl, O-(C1-C4)-alkyl or O-(C1-C4)-alkenyl and carbonyl;
- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl,

O-phenyl, amino, 1,2-methylenedioxy, carbonyl and phenyl;

- wherein R_1 is either hydrogen or U; X is either oxygen or CH-U, provided that if R_1 is hydrogen, then X is CH-U, or if X is oxygen then R_1 is U;

- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar (Ar as described above);

- wherein J is hydrogen or C1 or C2 alkyl or benzyl; K is (C1-C4)-straight or branched alkyl, benzyl or cyclohexylethyl; or wherein J and K may be taken together to form a 5-7 membered heterocyclic ring which may contain an oxygen (O), sulfur (S), SO or SO₂ substituted therein; and

- wherein n is 0-3.

26. The method of claim 25, further comprising co-administering an effective amount of a neurotrophic factor to promote neuronal regeneration selected from the group consisting of nerve growth factor, brain derived growth factor, glial derived growth factor, and neurotrophin-3.

27. The method of claim 25, wherein J and K are taken together to form a 5 membered heterocyclic ring.

28. The method of claim 27,

- wherein Z and R_2 are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-

straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns;

- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar; and

- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, and 1,2-methylenedioxy.

29. The method of claim 28, wherein

- if X is O, Y is O, NH or N-(C1-C4 alkyl), R₁ is C1-C6 straight or branched alkyl, C2-C6 straight or branched alkenyl, C5-C7 cycloalkyl or cycloalkenyl substituted with C1-C4 straight or branched alkyl or C2-C4 straight or branched alkenyl, and Ar is 1-naphthyl, 2-naphthyl, indolyl, 2-furyl, 3-furyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl

- then (CH₂)_nZ and R₂ taken together do not form:

1) substituted or unsubstituted indolyl, 2-furyl, 3-furyl, thiazolyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, phenyl

2) an alkyl or alkenyl chain with substituted or unsubstituted indolyl, 2-furyl, 3-furyl, thiazolyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, phenyl

3) an alkyl or alkenyl chain substituted with C5-C7 cycloalkyl.

30. The method of claim 25, wherein J and K are taken together to form a 6 membered heterocyclic ring.


31. The method of claim 30,

- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns;

- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar; and
- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, and 1,2-methylenedioxy.

32. The method of claim 31, wherein
- if n is 0, X is O, Y is O, R₁ is 1,1-dimethyl-1-propyl, and R₂ is 2-phenylethyl, then Z is not 1,1-dimethyl-1-prop-2-enyl, phenyl, cyclohexyl, or 3-(N,N-diallyl)benzamide; and
 - if n is 0, X is O, Y is O, R₁ is tert-butyl, and R₂ is 2-(3',4',5'-trimethoxyphenyl)ethyl, then Z is not phenyl.
33. A method for stimulating the growth of damaged peripheral nerves, comprising:


administering to said damaged peripheral nerves an effective amount of a compound having an affinity for FKBP-type immunophilins according to formula I



Formula I

or a pharmaceutically acceptable salt thereof,

- wherein Y is CH₂, O, NH, or N-(C1-C4 alkyl);
- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns, or



- wherein Q is hydrogen, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl;
- wherein T is Ar or substituted 5-7 membered cycloalkyl with substituents at

D5

positions 3 and 4 which are independently selected from the group consisting of hydrogen, hydroxyl, O-(C1-C4)-alkyl or O-(C1-C4)-alkenyl and carbonyl;

- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, 1,2-methylenedioxy, carbonyl and phenyl;

- wherein R₁ is either hydrogen or U; X is either oxygen or CH-U, provided that if R₁ is hydrogen, then X is CH-U, or if X is oxygen then R₁ is U;

- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar (Ar as described above);

- wherein J is hydrogen or C1 or C2 alkyl or benzyl; K is (C1-C4)-straight or branched alkyl, benzyl or cyclohexylethyl; or wherein J and K may be taken together to form a 5-7 membered heterocyclic ring which may contain an oxygen (O), sulfur (S), SO or SO₂ substituted therein; and

- wherein n is 0-3.

101

101

34. The method of claim 33, further comprising co-administering an effective amount of a neurotrophic factor to stimulate growth of the damaged peripheral nerve selected from the group consisting of nerve growth factor, brain derived growth factor, glial derived growth factor, and neurotrophin-3.

35. The method of claim 33, wherein J and K are taken together to form a 5 membered heterocyclic ring.

36. The method of claim 35,

- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns;

- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar; and

- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which

may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, and 1,2-methylenedioxy.

37. The method of claim 36, wherein

- if X is O, Y is O, NH or N-(C1-C4 alkyl), R₁ is C1-C6 straight or branched alkyl, C2-C6 straight or branched alkenyl, C5-C7 cycloalkyl or cycloalkenyl substituted with C1-C4 straight or branched alkyl or C2-C4 straight or branched alkenyl, and Ar is 1-naphthyl, 2-naphthyl, indolyl, 2-furyl, 3-furyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl

- then (CH₂)_nZ and R₂ taken together do not form:

- 1) substituted or unsubstituted indolyl, 2-furyl, 3-furyl, thiazolyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, phenyl
- 2) an alkyl or alkenyl chain with substituted or unsubstituted indolyl, 2-furyl, 3-furyl, thiazolyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, phenyl
- 3) an alkyl or alkenyl chain substituted with C5-C7 cycloalkyl.

38. The method of claim 33, wherein J and K are taken together to form a 6 membered heterocyclic ring.

39. The method of claim 38,


- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns;

- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar; and

- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, and 1,2-methylenedioxy.

40. The method of claim 39, wherein
- if n is 0, X is O, Y is O, R₁ is 1,1-dimethyl-1-propyl, and R₂ is 2-phenylethyl, then Z is not 1,1-dimethyl-1-prop-2-enyl, phenyl, cyclohexyl, or 3-(N,N-diallyl)benzamide; and
 - if n is 0, X is O, Y is O, R₁ is tert-butyl, and R₂ is 2-(3',4',5'-trimethoxyphenyl)ethyl, then Z is not phenyl.

41. A method for stimulating neurite outgrowth by a nerve cell, comprising:
- administering to said nerve cell an effective amount of compound having an affinity for FKBP-type immunophilins according to formula I




Formula I

or a pharmaceutically acceptable salt thereof,

- wherein Y is CH₂, O, NH, or N-(C1-C4 alkyl);
- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2

heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns, or



- wherein Q is hydrogen, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl;
- wherein T is Ar or substituted 5-7 membered cycloalkyl with substituents at positions 3 and 4 which are independently selected from the group consisting of hydrogen, hydroxyl, O-(C1-C4)-alkyl or O-(C1-C4)-alkenyl and carbonyl;
- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, 1,2-methylenedioxy, carbonyl and phenyl;
- wherein R₁ is either hydrogen or U; X is either oxygen or CH-U, provided that if R₁ is hydrogen, then X is CH-U, or if X is oxygen then R₁ is U;
- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkyl, (C5-C7)-cycloalkenyl substituted

with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar (Ar as described above);

- wherein J is hydrogen or C1 or C2 alkyl or benzyl; K is (C1-C4)-straight or branched alkyl, benzyl or cyclohexylethyl; or wherein J and K may be taken together to form a 5-7 membered heterocyclic ring which may contain an oxygen (O), sulfur (S), SO or SO₂ substituted therein; and

- wherein n is 0-3.

42. The method of claim 41, further comprising co-administering an effective amount of a neurotrophic factor to stimulating neurite outgrowth selected from the group consisting of nerve growth factor, brain derived growth factor, glial derived growth factor, and neurotrophin-3.

43. The method of claim 41, wherein J and K are taken together to form a 5 membered heterocyclic ring.

44. The method of claim 43,

- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns;

- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar; and

- wherein Ar is selected from the group consisting of monocyclic and bicyclic heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, and 1,2-methylenedioxy.

45. The method of claim 44, wherein

- if X is O, Y is O, NH or N-(C1-C4 alkyl), R₁ is C1-C6 straight or branched alkyl, C2-C6 straight or branched alkenyl, C5-C7 cycloalkyl or cycloalkenyl substituted with C1-C4 straight or branched alkyl or C2-C4 straight or branched alkenyl, and Ar is 1-naphthyl, 2-naphthyl, indolyl, 2-furyl, 3-furyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or phenyl

- then (CH₂)_nZ and R₂ taken together do not form:

- 1) substituted or unsubstituted indolyl, 2-furyl, 3-furyl, thiazolyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, phenyl
- 2) an alkyl or alkenyl chain with substituted or unsubstituted indolyl, 2-furyl, 3-furyl, thiazolyl, 2-thienyl, 3, thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, phenyl
- 3) an alkyl or alkenyl chain substituted with C5-C7 cycloalkyl.

46. The method of claim 41, wherein J and K are taken together to form a 6 membered heterocyclic ring.

47. The method of claim 46,

- wherein Z and R₂ are independently Ar, (C5-C7)-cycloalkyl substituted (C1-C6)-straight or branched alkyl or alkenyl, (C5-C7)-cycloalkenyl substituted (C1-C6)-straight or branched alkyl or alkenyl, or Ar substituted (C1-C6)-straight or branched alkyl or alkenyl, wherein in each case, one or two carbon atoms of the straight or branched alkyl or alkenyl groups may be substituted with 1-2 heteroatoms selected from the group consisting of oxygen, sulfur, SO and SO₂ in chemically reasonable substitution patterns;
- wherein U is hydrogen, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, (C5-C7)-cycloalkenyl substituted with (C1-C4)-straight or branched alkyl or (C1-C4)-straight or branched alkenyl, [(C1-C4)-alkyl or (C1-C4)-alkenyl]-Ar or Ar; and
- wherein Ar is selected from the group consisting of monocyclic and bicyclic

heterocyclic aromatic ring systems with individual ring sizes being 5 or 6 which may contain in either or both rings a total of 1-4 heteroatoms independently selected from oxygen, nitrogen and sulfur; wherein Ar may contain one to three substituents which are independently selected from the group consisting of hydrogen, halo, hydroxyl, hydroxymethyl, nitro, CF₃, trifluoromethoxy, (C1-C6)-straight or branched alkyl or (C1-C6)-straight or branched alkenyl, O-(C1-C4)-straight or branched alkyl or O-(C1-C4)-straight or branched alkenyl, O-benzyl, O-phenyl, amino, and 1,2-methylenedioxy.

48. The method of claim 47, wherein

-if n is 0, X is O, Y is O, R₁ is 1,1-dimethyl-1-propyl, and R₂ is 2-phenylethyl, then Z is not 1,1-dimethyl-1-prop-2-enyl, phenyl, cyclohexyl, or 3-(N,N-diallyl)benzamide; and

-if n is 0, X is O, Y is O, R₁ is tert-butyl, and R₂ is 2-(3',4',5'-trimethoxyphenyl)ethyl, then Z is not phenyl.